



## Garden Dust

8 oz dust

### **PRECAUTIONARY STATEMENTS: Hazard to Humans and Domestic Animals:**

**CAUTION:** Harmful if inhaled. Remove and wash contaminated clothing before reuse. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling this product and before eating, drinking, chewing gum or using tobacco.

#### **FIRST AID:**

##### **If inhaled:**

- Move person to fresh air.
- If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
- Call a poison control center or doctor for treatment advice.

##### **If on skin:**

- Take off contaminated clothing.
- Rinse skin immediately with water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

##### **If in eyes:**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor for treatment advice.

**Have the container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-858-7378 for emergency medical treatment advice.**

## **ENVIRONMENTAL HAZARDS**

Do not apply directly to water. Do not contaminate water when disposing of equipment washwater or rinsate. This product must not be applied within 300 feet of any habitats of endangered or threatened Lepidoptera (i.e. moths or butterflies.)

## **Directions for use:**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

## **Application Rates:**

Apply Safer® Brand Garden Dust when pests are first noticed and repeat every 7 to 10 days as necessary. Caterpillar egg hatch can occur over an extended time period. Monitor treated areas for the presence of newly hatched pests and re-treat if necessary. Thorough leaf coverage is necessary for effective control. After ingesting a treated portion of the leaf, caterpillars stop feeding within a few hours. Death occurs in a few days.

Apply to foliage of listed crops at a rate of 2 to 3 ounces per 50 feet of row planting.

## **Use on these crops:**

Broccoli	Sweet Potato	Melons
Cauliflower	Peppers	Spinach
Turnips	Lentils	Kale
Celery	Cucumbers	Turnip Greens
Cabbage	Squash	Mustard
Chinese Cabbage	Collard Lettuce	Brussels Sprouts
Kohlrabi	Beets	Sweet Corn
Tomatoes	Beans	Eggplant
Escarole	Endive	Potato
Carrots	Peas	Tobacco

## **To control these insects:**

Armyworms	Loopers	Tomato Fruitworm
Diamondback Moth	Melonworm	Corn Earworm
Green Cloverworm	Pickleworm	Bollworm
Hornworms	Mimosa Webworm	Tobacco Budworm
Imported Cabbageworm	Salt Marsh Caterpillar	

\*

\*

\*

## **Use on these fruits:**

Blackberries	Dewberries	Strawberries
Blueberries	Grapes	Currants
Raspberries		

**To control these insects:**

Armyworms	Grape Leafroller/Leafroller	Loopers
Grapeleaf Skeletonizer	Hornworms	Salt Marsh Caterpillar

\* \* \*

**Use on Lawns to Control this Insect:**

Sod Webworm

\* \* \*

**Use on Flowers and Ornamentals to Control these Insects:**

Loopers	Tobacco Budworm	Omnivorous Loopers
Omnivorous Leaf Roller	Diamondback Moth	Armyworms
Ello Moth (Hornworm)	Io Moth	Oleander Moth

\* \* \*

**STORAGE AND DISPOSAL:**

Do not contaminate water, food or feed by storage and disposal.

**Storage:** Store in a cool dry area away from heat or open flame and inaccessible to children and animals.

**Disposal:**

**If empty:** Do not reuse this container. Place in the trash or offer for recycling if available.

**If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.